Application No.: 10/532,160

Office Action dated: September 18, 2008 Reply to Office Action Dated: January 21, 2009 Attorney Docket No. 13779-21 Page 2 of 7

Amendments to the claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended): Process for the preparation of a compound of general formula (I):

$$X$$
 NH_2
 X
 CF_3
 (I)

in which X represents a halogen atom, by reaction of para-trifluoromethylaniline of formula (II):

with a dihalogen X_2 ,

the two compounds being introduced simultaneously into a polar aprotic solvent in a dihalogen/compound (II) molar ratio ranging from 1.9 to 2.5 and at a temperature ranging from 100 to 300°C, without the addition of hydrofluoric acid.

- 2. (Previously presented): Process according to Claim 1, characterised in that the compound of formula (I) is 2,6-dichloro-para-trifluoromethylaniline.
- 3. (Previously presented): Process according to Claim 1, wherein the solvent used is a chlorinated aliphatic solvent.

Application No.: 10/532,160
Office Action dated: September 18, 2008

Office Action dated: September 18, 2008 Reply to Office Action Dated: January 21, 2009

4. (Previously presented): Process according to Claim 3, characterised in that the

Attorney Docket No. 13779-21

solvent used is dichloroethane.

5. (Previously presented): Process according to Claim 1, wherein the solvent used is

a chlorinated aromatic solvent.

6. (Previously presented): Process according to Claim 5, characterised in that the

solvent used is monochlorobenzene.

7. (Previously presented): Process according to Claim 1, wherein, the reactants are

introduced in a dihalogen/compound (II) molar ratio ranging from 2 to 2.05.

8. (Previously presented): Process according to Claim 1, wherein the temperature of

the reaction medium is chosen as ranging from 100 to 130°C.

9. (Previously presented): Process according to Claim 8, characterised in that the

temperature of the reaction medium is chosen as ranging from 105 to 115°C.

10. (Previously presented): Process according to Claim 2, characterized in that the

reactants are introduced into monochlorobenzene in a dichlorine/compound (II) molar ratio

ranging from 1.85 to 2.05, at a temperature ranging from 105 to 115°C.

11. (Previously presented): Process according to Claim 2, wherein the solvent used is

a chlorinated aliphatic solvent.

12. (Previously presented): Process according to Claim 2, wherein the solvent used is

a chlorinated aromatic solvent.